

Tagung der Gesellschaft für Immunologie
Society of Immunology L'Association d'Immunologie
Berlin Oct 5-8, 1983
XVth Meeting of the Society of Immunology · 323

Institut für Immunologie und Serologie, Universität Heidelberg, Heidelberg, FRG

143. Immunological and functional properties of two monoclonal antibodies against human C5

T. MOONCKARNDT, A. DESSAUER, H. BRAUCH, G. M. HÄNSCH, and E. W. RAUTERBERG

Two monoclonal antibodies designated 4C6 and 3B2 were produced by immunizing Balb/c 4C6 or DBA₂ mice (3B2) repeatedly with human C5 purified according to DESSAUER et al. (Immunobiol., in press) and following the fusion of the mouse spleen cells with NS-1 (4C6) or NS-3 Ag8.653 (3B2) lymphoma cells respectively. Hybridomas have been selected according to the results of ELISA and RIA measurements with insolubilized human C5. They were cloned and IgG fractions of ascites were isolated by 45% saturated ammonium sulfate precipitation and DEAE ion exchange chromatography. These fractions were investigated by analytical and functional assays.

The antibodies (ab) in one of the lines (4C6) reacted with a 200 kD protein after SDS-PAGE and immunoblotting (IB) of unreduced human C5, whereas line 3B2 reacted with a 60 kD peptide chain, which is suspected to represent a split product of C5, since its proportion increased with time during the storage of C5. Line 4C6 failed to react with reduced C5 in the IB analysis, in contrast line 3B2 reacted even after reduction of C5 with the 60 kD band. Functionally this antibody (3B2) inhibited the ³H serotonin release from guinea pig platelets induced by hog C5a and surprisingly also the lysis of chicken erythrocytes by C5 + C7, C8, C9 and the lysis of EAC1-5 by C6-9 or EAC1-6 by C7-9, but failed to inhibit the interaction of C5 with EAC1423 in the presence of C6-9 in excess. The other antibody (4C6), after preincubation with C5 strongly inhibited the C5 dependent lysis of erythrocytes when studied with EAC1423 + C6-9 and also diminished the lysis of ChE by C56 and C7-9, but failed to react with EAC1-6.

Our findings suggest that one of the monoclonal ab (4C6) detects an epitope on the C5 molecule related to its reaction with target membranes. This epitope seems to be hidden or without further functional role on EAC1-5 or EAC1-6.

The other monoclonal ab (3B2) displaying functional inhibitory activities puzzles by its functional inhibitory effect on C5a dependent serotonin release on one side and its reaction with cell-bound C5 or free and bound C5b6. Further studies are needed to clarify, whether these contradictory results are due to the possibility, that C5a is produced by cleavage but remains noncovalently bound to C5b.

Max-Planck-Institut für Immunbiologie, Freiburg i. Breisgau, FRG

144. Potentiation of antibody-dependent cellular cytotoxicity and chemiluminescence in human neutrophils by platelet activating factor

H. MOSSMANN, U. BAMBERGER, B. A. VELEV, D. K. HAMMER

Human neutrophils (PMN) purified by elutriation from blood of healthy volunteers were tested in the presence of platelet activating factor (PAF) for both antibody-dependent cellular cytotoxicity (ADCC) against antibody coated erythrocytes and the oxidative response as

250 ug/ml
100 ug/ml

ology

r splenic T-cells. Increased Ig as measured by secretion only as shown by studying de novo Ig cellular and secreted Ig. The system was used to increase in Ig synthesis. Marrack's group is identical to γ -interferon. Using back, that γ -interferon per se is totally ineffective. However, a combination of cloned γ -interferon B cell differentiation factors lead to some of doses of BCGs were used. Kinetic studies "priming effect", making cells more receptive on factors.

des Krebsforschungszentrum, 6900 Heidelberg
3 Heidelberg, FRG

Idiotypes against a human B cell
for monitoring and therapy

(12), and G. J. HAMMERLING

thomas were found to be of clonal origin. In proliferation of a single B lymphocyte clone, i.e. restricted to the expression of identical ygs (idiotypes). Therefore, the unique idiotype is tumor-specific marker against which anti-

as the immunoglobulin molecule is expressed mounts. In order to rescue immunoglobulin used. Peripheral blood mononuclear cells from were fused with mouse myeloma cells. Line 273 secreted human immunoglobulin. tumor cell isotype (γ , λ). One representative several times, and propagated in bulk culture. ant was purified by affinity chromatography. immunoglobulin and hybridomas were generated. At least 3 different monoclonal antibodies identified. Specificity of these antibodies was tion assay. The 3 antibodies react exclusively civity with unrelated immunoglobulins was re actually individual-specific.

red for quantitative detection of idiotype re serum during the course of disease. The residual tumor cells after chemotherapy. The for autologous bone marrow transplantation

Immunobiology
165 (1983)

3B2